

SUSY WITHOUT R -PARITY: SYMMETRY BREAKING AND LSP-PHENOMENOLOGY

Ralf Hempfling
Davis Institute for High Energy Physics
Davis, CA 95616, USA
 E-mail: hempf@bethe.ucdavis.edu

ABSTRACT

We have studied the predictions for the LSP decay within the framework of a radiatively broken unified supergravity model without R -parity. Assuming that Higgs/slepton mixing is the only source of R -parity breaking and responsible for the observed neutrino oscillations we obtain predictions for the LSP life-time and branching fractions.

1. Motivation

Supersymmetry¹ is presently the most popular attempt to solve the hierarchy problem of the standard model (SM). Here, the cancellation of quadratic divergences is guaranteed and, hence, any mass scale is stable under radiative corrections. The most economical candidate for a realistic model is the minimal supersymmetric extension of the SM (MSSM). In the SM baryon and lepton number are protected by an accidental symmetry (*i.e.* there is no gauge and Lorentz invariant term of dimension 4 or less that violates B or L via perturbative effects). This no longer holds in the MSSM due to the existence of superpartners. One way to assure Baryon and Lepton number conservation (and hence the stability of the proton) is to impose by hand a discrete, multiplicative symmetry called R -parity², $R_p = (-1)^{2S+3B+L}$, where S , B and L are the spin, baryon and lepton numbers, respectively. Aside from the long proton life-time, R_p conserving models have the very attractive feature that the lightest supersymmetric particle (LSP) is stable and a good cold dark matter candidate³. On the other hand, there is strong experimental evidence for neutrino oscillations^{4,5,6} which can be accounted for if R_p is broken⁷.

In this paper, we will investigate the LSP life-time in a SUSY-GUT scenario where R_p is broken explicitly via dimension 2 terms^{7,8,9,10}. We have discussed this model detail in ref. 7 where the emphasis was on neutrino phenomenology in the frame-work of radiative electro-weak spontaneous symmetry breaking (REWSSB)¹¹. Here, we are particularly interested in the implications for high energy collider phenomenology. We will focus attention of the case that the LSP is a neutralino. This is the most interesting case, since it occurs naturally over most of the SUSY parameter space. Note, however, that in models with broken R_p there is no theoretical/cosmological prejudice concerning the color or electric charge of the LSP. The only requirement is that the LSP life-time is sufficiently short ($\tau_{LSP} \lesssim 1$ sec) so as not to disturb big

bang nucleosynthesis¹² or sufficiently long [$\tau_{LSP} \gtrsim 10^{24} \text{ sec}/B(LSP \rightarrow X\nu_e)$] so as not to lead to an unacceptable distortion of the cosmic microwave background¹³.

The most general gauge invariant superpotential can be written as

$$W = \frac{1}{2}y_{IJk}^L \hat{L}_I \hat{L}_J \hat{E}_k^c + y_{Ijk}^D \hat{L}_I \hat{Q}_j \hat{D}_k^c - y_{jk}^U \hat{H} \hat{Q}_j \hat{U}_k^c - \mu_I \hat{L}_I \hat{H} + \frac{1}{2}\bar{y}_{ijk}^D \hat{D}_i^c \hat{D}_j^c \hat{U}_k^c, \quad (1)$$

where the supermultiplets are denoted by a hat. The left-handed lepton supermultiplets are denoted by $\hat{L}_i$ ($i = 1, 2, 3$) and the Higgs superfield coupling to the down-type quarks is denoted by $\hat{L}_0$.

Let us first determine the meaning of the various terms of eq. 1. Here, y_{0jk}^L , y_{0jk}^D and y_{jk}^U denote the lepton, down-type and up-type Yukawa couplings, respectively, and μ_0 is the Higgs mass parameter. However, in contrast to the SM the MSSM allows for renormalizable baryon [lepton] number violating interactions $\bar{y}_{ijk}^D$ [y_{ijk}^L and μ_i]. These couplings are constrained from above by experiment. The most model independent constraints can be obtained from collider experiments¹⁴ or neutrino physics¹⁵. It turns out that the individual lepton and baryon number violating couplings only have to be smaller than $O(10^{-3} \sim \text{few} \times 10^{-1})$ with the exception of $\bar{y}_{121}^D \lesssim 10^{-7}$ from heavy nuclei decay¹⁶. Thus, the R_p violating couplings need not be much more suppressed than the lepton and baryon number preserving Yukawa couplings. (remember that *e.g.*, $y_{011}^D \simeq 3 \times 10^{-5}/\cos\beta$). Somewhat stronger but more model dependent constraints can be derived from cosmology¹⁷.

However, the experimental exclusion area can be strongly enhanced by imposing theoretical constraints: in the minimal SU(5) SUSY-GUT model, the right-handed leptons, the right-handed up-type quarks and the left-handed quarks are embedded in a 10-dim representation, $\mathbf{10}_i = E_i^c \oplus U_i^c \oplus Q_i$. The right-handed down-type quarks and the left-handed leptons are embedded in a 5-dim representation, $\bar{\mathbf{5}}_i = D_i \oplus L_i$. The two Higgs doublets are embedded together with two proton decay mediating colored triplets, T and D_0 , in 5-dim representations, $\bar{\mathbf{5}}_0 = D_0 \oplus L_0$ and $\mathbf{5} = T \oplus H$. Hence, both the lepton and the baryon number violating interactions arise from the term

$$W_{\text{GUT}} = \frac{1}{2}y_{ijk} \bar{\mathbf{5}}_i \bar{\mathbf{5}}_j \mathbf{10}_k, \quad (2)$$

where the boundary conditions at the GUT-scale, M_{GUT} , are given by $y_{ijk}^L = y_{ikj}^D = \bar{y}_{ijk}^D = y_{ijk}$. These relations, which predict the down-type quark masses correctly to within a factor of 3, are expected to also hold at a comparable level for the R_p violating couplings. Thus, in general the baryon and lepton number violating couplings are correlated in SUSY GUT models. This leads to very strong constraints on any y_{ijk} from proton stability which are much stronger than any constraint on individual Yukawa couplings¹⁸. However, it does not place any constraints on the coefficients of of dimension 2 terms, μ_i , which are the subject of this paper.

The outline of our paper is as follows: in section 2 we present the neutrino and sparticle spectrum obtained from REWSSB including μ_i . In section 3 we present




Fig. 1. Contours of constant m_0 , B_0 and $\sin \theta'_1{}^{\text{app}}/\sin \theta'_1$ in the $\tan \beta$ - $m_{1/2}$ plane. We set $A_0 = 0$ and $\mu = 2.5m_{1/2}$.

the numerical analysis of the LSP life-time, τ_{LSP} , and LSP branching fractions. Our conclusions are presented in section 4.

2. Radiative Electro-Weak Symmetry Breaking

Without any assumptions based on theoretical prejudice there are many models with different SUSY particle spectra and vastly different phenomenology. Thus, it has become standard to derive the low energy particle spectrum from minimal supergravity model with only four independent parameters: the universal scalar mass parameter, m_0 , the universal gaugino mass parameter, $m_{1/2}$, and the universal A (B) parameter multiplying the tri-linear (bi-linear) terms in the superpotential [eq. 1]. This approach is supported by the observation that the absence of FCNC implies a high mass-degeneracy of all scalars with the same gauge quantum numbers (with the possible exceptions of the Higgs mass parameters).

First, we have to minimize the Higgs potential given by

$$\begin{aligned}
 V = & (\mu^2 + m_H^2)H^\dagger H + (\mu_I \mu_J + m_{L_{IJ}}^2)\tilde{L}_I^\dagger \tilde{L}_J - B_{IJ}\mu_I (\tilde{L}_J H + \text{H.c.}) \\
 & + \frac{g^2 + g'^2}{8} (H^\dagger H - \tilde{L}_I^\dagger \tilde{L}_I)^2 + \frac{g^2}{2} |H^\dagger \tilde{L}_I|^2,
 \end{aligned} \tag{3}$$

where the low energy soft SUSY breaking parameters are obtained by renormalization group evolution below M_{GUT} in the standard fashion. In order to stay as close to the notation of the MSSM as possible we follow our notation of ref. 7

$$\bar{v} \equiv \frac{\langle H^0 \rangle}{\sqrt{2}}, \quad v_I \equiv \frac{\langle \tilde{L}_I^0 \rangle}{\sqrt{2}}, \quad v \equiv \sqrt{v_I v_I}, \quad \text{and} \quad \tan \beta \equiv \bar{v}/v, \tag{4}$$

and we parameterize the VEVs in terms of spherical coordinates

$$\tan \theta'_3 = \frac{v_3}{v_2}, \quad \tan \theta'_2 = \frac{v_2}{v_1 \cos \theta_3}, \quad \tan \theta'_1 = \frac{v_1}{v_0 \cos \theta_2}. \tag{5}$$




Fig. 2. Contours of constant (a) $m_{\tilde{t}_1}$, (b) $m_{\tilde{b}_1}$, (c) m_{A^0} (tree-level) and (d) m_{h^0} (incl. 1-loop radiative corrections) in the $\tan\beta$ - $m_{1/2}$ plane. The other SUGRA parameters are as in fig. 1.

Analogously, it is convenient to parameterize the R_p breaking mass parameters in terms of three mixing angles

$$\tan\theta_3 = \frac{\mu_3}{\mu_2}, \quad \tan\theta_2 = \frac{\mu_2}{\mu_1 \cos\theta_3}, \quad \tan\theta_1 = \frac{\mu_1}{\mu_0 \cos\theta_2}, \quad (6)$$

and $\mu \equiv \sqrt{\mu_I \mu_I}$. The potential in eq. 3 is minimized by an iterative procedure using the analytic solution for $\tan\theta_1 = 0$ as our initial values. This procedure also works surprisingly well for $\tan\theta_1 > 1$. For small R_p violating parameters we can also obtain very reliable analytic expressions in the basis where y_{ij}^L is diagonal by expanding in powers of μ_i/μ_0

$$\sin 2\beta = \frac{2B_{00}\mu_0}{m_{L00}^2 + m_H^2 + 2\mu_0^2}, \quad (7)$$

$$\tan^2 \beta = \frac{m_{L00}^2 + \mu_0^2 + \frac{1}{2}m_z^2}{m_H^2 + \mu_0^2 + \frac{1}{2}m_z^2}, \quad (8)$$

$$\frac{v_i}{v_0} = \mu_i \frac{B_{(ii)} \tan\beta - \mu_0}{m_{L(ii)}^2 + \frac{1}{2}m_z^2 \cos 2\beta}, \quad (9)$$

with the convention that indices in braces are not summed over. In general, one fixes the GUT input parameters are m_0 , $m_{1/2}$ and A_0 . B_{00} and μ are obtained by solving




Fig. 3. Contours of constant (a) $m_{\chi_1^0}$, (b) $m_{\chi_1^\pm}$, in the $\tan\beta$ - $m_{1/2}$ plane. The other SUGRA parameters are as in fig. 1.

eq. 7 and 8 while keeping $\tan\beta$ and v fixed. Here, we find it convenient to fix the fermionic spectrum given by μ rather than the scalar spectrum determined by m_0 .

In fig. 1(a) and (b) we present contours of constant GUT parameters m_0 and B_0 in the $\tan\beta$ - $m_{1/2}$ plane. and in fig. 1(c) we show how well the approximation works for the minimization of the potential (eq. 9). We see that the deviation of the approximation obtained from eq. 9 denoted by $\sin\theta_1^{\text{app}}$ from the true minimum obtained by numerical methods and denoted by $\sin\theta_1'$ is quite small as long as $\tan\beta = O(1 \sim 10)$. However, it breaks down for $\tan\beta \gtrsim 40$.

2.1. Sparticle Spectrum

From LEP experiments we know that there are no charged superpartners with mass below $m_z/2$. Furthermore, we can deduce a similar constraint on the lightest neutralino which, in our model, is instable. In fig. 2 and 3 we present contours of some relevant scalar and fermionic superpartner masses in the $\tan\beta$ - $m_{1/2}$ plane. We have chosen $A_0 = 0$ and $\mu = 2.5m_{1/2}$. The value of m_0 is obtained by imposing REWSSB [see fig. 1(a)]. We see that the only relevant constraint arises from the experimental lower on the lightest neutralino mass denoted by M_{LSP} .

2.2. Neutrino Spectrum

The LSP phenomenology of the model under investigation here is governed to a very good approximation by only one parameter, $\tan\theta_1$. This parameter also determines the neutrino masses whose upper limits are given by

$$\begin{aligned}
 m_{\nu_e} &\leq 4.35 \text{ eV}, m_{\nu_\mu} \leq 160 \text{ keV}, m_{\nu_\tau} \leq 23 \text{ MeV}, & \text{Collider-experiment}^{19} \\
 \sum_{x=e,\mu,\tau} m_{\nu_x} &\leq (10 \sim 100) \text{ eV}, & \text{Cosmology}^{20} \quad (10)
 \end{aligned}$$




Fig. 4. Contours of constant (a) $m_{\nu_\mu}^2 - m_{\nu_e}^2$ (b) $\sin^2 2\theta_{\tau\nu_\mu}$ in the $\tan\beta$ - M_{LSP} plane. We set $A_0 = 0$ and $\mu = 2.5m_{1/2}$ and $m_{1/2}$ is replaced in favor of M_{LSP} .

While there is no direct evidence for non-zero neutrino masses, there is strong experimental evidence for neutrino oscillations which imply that the three neutrinos are non mass-degenerate. In this work, we will take the view that our R_p violating terms are the only source of neutrino masses and, therefore, should account for all the existing neutrino mixing effects. Since not all experimental indication for neutrino oscillations appear compatible with each other or have the same statistical significance a selection has to be taken.^a The solar neutrino puzzle⁴ may be the most compelling evidence for neutrino mixing. However, the effect appears to involve only the first two neutrino flavors neither of which is likely to be the heaviest neutrino both on theoretical and experimental grounds. Thus, the solar neutrino puzzle is not well suited for a determination of $\tan\theta_1$. Instead we choose to solve the atmospheric neutrino problem⁵. This requires that we fix $\tan\theta_1$ such that $m_{\nu_\tau}^2 - m_{\nu_\mu}^2 = 10^{-2} \text{ eV}^2$ and we set $\tan\theta_2 = 1$ (we use a small value for $\tan\theta_3 = 0.045$ in order to solve the solar neutrino puzzle via the MSW-effect²²; this angle will turn out to be quite irrelevant otherwise). Over most of the parameter space under consideration here this implies $m_{\nu_\mu} \ll m_{\nu_\tau} \simeq 0.1 \text{ eV}$. It is then straightforward to obtain lower limits on τ_{LSP} from upper limits on m_{ν_τ} [eq. 10] by simple scaling arguments.

^aIt has recently been suggested that three neutrino flavor are enough to accomodate all three indication for neutrino oscillation²¹. However, for the sake of generality we will be more conservative.




Fig. 5. The LSP-width, Γ_{LSP} (divided by $\tan^2 \theta_1$) and the mass difference $m_{\nu_\tau}^2 - m_{\nu_\mu}^2$ (divided by $\tan^4 \theta_1$) as a function of $\tan \theta_1$, $\tan \theta_2$ and $\tan \theta_3$.

In fig. 4 we present contours of constant values for $m_{\nu_\mu}^2 - m_{\nu_e}^2$ and $\sin^2 2\theta_{\mu\nu\tau}$. We fix $\mu = \pm 2.5 m_{1/2}$ and $A_0 = 0$. We see that for positive values of μ the mass difference $m_{\nu_\mu}^2 - m_{\nu_e}^2 = O(10^{-9} \text{ eV}^2)$ is very small and quite compatible with long wave oscillation (LWO)²³ solution to the solar neutrino problem [fig. 4 does not change if we set $\tan \theta_3 = O(1)$]. For $\mu < 0$ there is a sizable region where $m_{\nu_\mu}^2 - m_{\nu_e}^2 = O(10^{-5} \text{ eV}^2)$ as required by the MSW explanation of the solar neutrino deficiency⁴. These results were first presented in ref. 7^b

3. LSP Phenomenology

In this section, we will discuss the decay properties of the LSP. The main interest from the point of view of collider phenomenology is the question whether the LSP decays inside the detector (else the analysis is equivalent to the case of unbroken R_p).

Since the magnitude of R_p violation is parameterized by an a priori free parameter $\tan \theta_1$, we cannot determine τ_{LSP} . The situation changes if we relate $\tan \theta_1$ to the neutrino masses. In the first part of this section we will present the prediction for τ_{LSP} assuming the atmospheric neutrino puzzle is a result of R_p violation. [This prediction can be turned into a lower limit by imposing any of the upper limits on m_{ν_τ} of eq. 10.] In the second part of this section, we will discuss the branching fractions of the LSP which is independent of $\tan \theta_1$ and, hence, also of any assumption about neutrino masses.

3.1. LSP life-time

As pointed out in ref. 8 in models without R_p the neutrinos and neutralinos are

^bNote that fig. 7(a), fig. 8(a) and fig. 9 are mislabeled in ref. 7. The region with $\mu > 0$ and the region with $\mu < 0$ should be interchanged.




Fig. 6. Contours of constant $\tan \theta_1$ and $\tau(\chi_1^0)$ in the $\tan \beta$ - M_{LSP} plane for $\mu = \pm 2.5m_{1/2}$. The other SUGRA parameters are as in fig. 4.

indistinguishable. As a result, the formulæ for the neutralino radiative decay in the MSSM ²⁴ and the decay into three fermions ²⁵ can be directly generalized to our model. However, we do have to include the effects of Yukawa couplings which were neglected for R_p preserving three-body decays ²⁵. Our complete set of formulæ will be presented elsewhere²⁶. Here, we simply present our numerical results.

In fig. 5 we have plotted Γ_{LSP} vs. $\tan \theta_1$. We find that there are simple scaling relations if R_p violation is sufficiently small (say $\tan \theta_1 \lesssim 0.1$):

$$\Gamma_{LSP}, m_{\nu_x} \propto \tan^2 \theta_1 \quad (x = e, \mu, \tau). \quad (11)$$

The parameters $\tan \theta_2$ and $\tan \theta_3$ which govern the neutrino oscillations have only a small impact on the LSP properties. In fig. 6 we present contours of $\tan \theta_1$ (fixed by imposing $m_{\nu_\tau}^2 - m_{\nu_e}^2 = 10^{-2} \text{ eV}^2$) and constant τ_{LSP} in the $\tan \beta$ - M_{LSP} plane for $\mu = \pm 2.5m_{1/2}$. We find that that required range of the R_p violation is $\tan \theta_1 = 10^{-x}$ ($x = 2 \sim 5$) with the upper (lower) limit corresponding to small (large) values of $\tan \beta$. The corresponding range of the LSP life-time is $c\tau_{LSP} = 1 \text{ m} \sim 0.1 \text{ mm}$ (for $M_{LSP} = 40 \sim 160 \text{ GeV}$) and can easily be determined at forthcoming collider experiments. Furthermore, we need larger R_p violation (for fixed m_{ν_τ}) for $\mu < 0$ due to cancellation among tree-level and one-loop contributions. For $\tan \beta > 30 \sim 40$ the one-loop contribution even dominates over the tree-level result. For $M_{LSP} \gtrsim m_z$ the




Fig. 7. A comparison of the total LSP-width with the LSP-width due to two-body decays. In (a) we show both sets of curves as a function of M_{LSP} . In (b) we show the difference of total width minus two-body decays normalized to Γ_{LSP} .




Fig. 8. The effects of non-universal boundary conditions at M_{GUT} on (a) τ_{LSP} , (b) $\tan\theta_1$ and (c) $m_{\nu_\mu}^2 - m_{\nu_e}^2$ for four different values of $\tan\beta$. We set $m_{1/2} = 120$ GeV, $\mu = 300$ GeV and $A_0 = 0$.

two-body decays dominate and differ from the full width only by a few % [fig. 7]. (A partial analysis of the two-body decays has been performed previously¹⁰.)

So far we have assumed exact universality at M_{GUT} . However, it has been pointed out in ref. 27 that the evolution from the Planck scale M_P to M_{GUT} can already have a significant impact on the sparticle spectrum. This is particularly important in SO(10) based models where the Higgs and slepton universality is violated via gaugino effects, since the Higgs (slepton) fields belong to a 10-dim (16-dim) representation (remember: below M_{GUT} non-universal effects arise only from Yukawa couplings while the non-universal effects above M_{GUT} arise from the gauge couplings). We




Fig. 9. Different LSP-branching fractions as functions of M_{LSP} for five values of $\tan\beta$. We set $A_0 = 0$ and $\mu = 2.5m_{1/2}$

can accommodate this effect by modifying the boundary conditions at M_{GUT}

$$m_H^2(M_{GUT}) = m_{L_0}^2(M_{GUT}) = m_0^2 + R_H m_{1/2}^2, \quad (12)$$

where we typically expect $R_H = 9\alpha_{GUT}/(4\pi)\ln(M_{GUT}/M_P) \simeq -0.1$.

In fig. 8 we see that the effect of non-universal terms is very significant (small) for small (large) values of $\tan\beta$ were the down-type Yukawa couplings are small (large).

3.2. LSP Branching Fraction

So far we have used results from neutrino physics in order to eliminate the R_p breaking parameters $\tan\theta_i$ ($i = 1, 2, 3$). However, to a good approximation this dependence drops out if we consider the branching fractions. In fig. 9 we present the branching fractions as a function of M_{LSP} for eight different channels. The dominant decay mode is into quarks [(a) and (b) are first two generations only; (e) is the third generation] with a strong enhancement into $b\bar{b}$ (e) for small $\tan\beta$. Invisible decay modes (c) are typically below 10% and the radiative decay (d) is insignificant. The leptonic decays into $\tau^+\tau^-$ (f), $\ell^+\ell^-$ ($\ell = e, \mu$) (g) and $\ell^\pm\tau^\mp$ (h) is typically O(10%). For $M_{LSP} \gtrsim 100$ GeV the situation becomes much more transparent by considering the two-body decays [fig. 10]. Here, there are only three relevant channels with $B(LSP \rightarrow W^\pm\tau^\mp) \simeq 0.5$ and $B(LSP \rightarrow Z^0\nu), B(LSP \rightarrow h^0\nu) \simeq 0.25$.




Fig. 10. Branching fractions of different two-body decays as functions of M_{LSP} for five values of $\tan\beta$. SUGRA parameters are same as in fig. 9.

4. Conclusions

We have investigated the LSP phenomenology in supersymmetric models without lepton number conservation. In any model of this kind, lepton number is violated spontaneously via sneutrino VEVs as well as explicitly. Both effects are of the same order and have to be studied consistently. Assuming that Higgs-sneutrino mixing is responsible for the observed neutrino oscillations we find that the LSP decays inside the detector. The life-time can be determined over a large region of the SUSY parameter space. The branching fractions for all relevant decay modes are we presented.

5. Acknowledgements

This work was supported in parts by the DOE under Grants No. DE-FG03-91-ER40674 and by the Davis Institute for High Energy Physics.

6. References

1. for a review, see, *e.g.*, H.P. Nilles, *Phys. Rep.* **110**, 1 (1984); H.E. Haber and G.L. Kane, *Phys. Rep.* **117**, 75 (1985).
2. G. Farrar and P. Fayet, *Phys. Lett.* **76B**, 575 (1978); N. Sakai and T. Yanagida, *Nucl. Phys.* **B197**, 533 (1982).
3. J. Ellis *et al.*, *Nucl. Phys.* **B238**, 453 (1984).
4. R. Davis *et al.*, *Phys. Rev. Lett.* **20**, 1205 (1968); K.S. Hirata *et al.*, *Phys. Rev. Lett.* **65**, 1297 (1990); S.A. Bludman *et al.*, *Phys. Rev.* **D47**, 2220 (1993); P. Anselmann *et al.*, *Phys. Lett.* **B327**, 377 (1994).

5. K.S. Hirata *et al.*, *Phys. Lett.* **B280**, 146 (1992). R. Becker-Szendy *et al.*, *Phys. Rev. Lett.* **69**, 1010 (1992); *Phys. Rev.* **D46**, 1992 (3720). Y. Fukuda *et al.*, *Phys. Lett.* **B335**, 237 (1994).
6. C. Athanassopoulos *et al.*, *Phys. Rev.* **D75**, 2650 (1995); LA-UR-96-1326.
7. R. Hempfling, *Nucl. Phys.* **B478**, 3 (1996).
8. L.J. Hall and M. Suzuki, *Nucl. Phys.* **B231**, 419 (1984).
9. T. Banks, Y. Grossman, E. Nardi and Y. Nir, *Phys. Rev.* **D52**, 5319 (1996); M. Nowakowski and A. Pilaftsis, *Nucl. Phys.* **B461**, 19 (1996); B. de Carlos and P.L. White, *Phys. Rev.* **D54**, 3427 (1996); H-P. Nilles and N. Polonsky, TUM-HEP-245-96 (May 1996).
10. S. Roy and B. Mukhopadhyaya, MPI-PHY/96-37, hep-ph/9612447.
11. J. Ellis, J.S. Hagelin, D.V. Nanopoulos and K. Tamvakis, *Phys. Lett.* **B125**, 275 (1983); G.F. Giudice and G. Ridolfi, *Z. Phys.* **C41**, 447 (1988).
12. for a review see: S. Sarkar, *Rep. Prog. Phys.* **59**, 1493 (1996).
13. M.V. Diwan, contribution to 1996 DPF/DPB Summer Study on *New Directions for High-energy Physics*, Snowmass, CO.
14. V. Barger, G.F. Giudice and T. Han, *Phys. Rev.* **D40**, 2987 (1989); H. Dreiner and G.G. Ross, *Nucl. Phys.* **B365**, 597 (1991); G. Bhattacharyya, D. Choudhury and K. Sridhar, *Phys. Lett.* **B355**, 193 (1995).
15. K. Enqvist, A. Masiero and A. Riotto, *Nucl. Phys.* **B373**, 95 (1992). J.C. Romao and J.W.F. Valle, *Nucl. Phys.* **B381**, 87 (1992).
16. C.E. Carlson, P. Roy and M. Sher, *Phys. Lett.* **B357**, 99 (1995).
17. B.A. Campbell, S. Davidson, J. Ellis and K.A. Olive, *Phys. Lett.* **B256**, 457 (1991); H. Dreiner and G.G. Ross, *Nucl. Phys.* **B410**, 88 (1993).
18. A.Yu Smirnov and F. Vissani, *Nucl. Phys.* **B460**, 37 (1996).
19. I.A. Belesev *et al.*, *Phys. Lett.* **B350**, 263 (1995); K. Assamagan *et al.*, *Phys. Lett.* **B335**, 231 (1994); D. Buskulic *et al.*, *Phys. Lett.* **B349**, 585 (1995).
20. for a review, see, e.g. E.W. Kolb and M.S. Turner, *The Early Universe*, (Addison-Wesley, Redwood City, CA, 1990).
21. A. Acker and S. Pakvasa, hep-ph/9611423
22. L. Wolfenstein, *Phys. Rev.* **D17**, 2369 (1978); *ibid.* **20**, 2634 (1979); S.P. Mickheyev and A. Yu Smirnov, *Yad. Fiz.* **42**, 1441 (1985) [*Sov. J. Nucl. Phys.* **42**, 913 (1986)].
23. V. Gribov and B. Pontecorvo, *Phys. Lett.* **B28**, 493 (1969); V. Barger, R.J.N. Phillips and K. Whisnant, *Phys. Rev.* **D24**, 538 (1981); *Phys. Rev. Lett.* **69**, 3135 (1992).
24. H.E. Haber and D. Wyler, *Nucl. Phys.* **B323**, 267 (1989).
25. A. Bartl, H. Fraas and W. Majerotto, *Z. Phys.* **C41**, 475 (1988).
26. H.E. Haber and R. Hempfling, in preparation.
27. N. Polonsky and A. Pomerol, *Phys. Rev. Lett.* **73**, 2292 (1994).

